

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**

**THIS PAGE BLANK (USPTO)**

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>7</sup> :</b> <b>A61K 38/00</b>	<b>A2</b>	<b>(11) International Publication Number:</b> <b>WO 00/12110</b> <b>(43) International Publication Date:</b> 9 March 2000 (09.03.00)
<b>(21) International Application Number:</b> PCT/CA99/00787 <b>(22) International Filing Date:</b> 25 August 1999 (25.08.99) <b>(30) Priority Data:</b> 60/098,178 26 August 1998 (26.08.98) US <b>(71) Applicant (for all designated States except US):</b> QUEEN'S UNIVERSITY AT KINGSTON [CA/CA]; Kingston, Ontario K7L 3N6 (CA). <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> ADAMS, Michael, A. [CA/CA]; 341 Renda Street, Kingston, Ontario K7M 5X9 (CA). HEATON, Jeremy, P., W. [CA/CA]; 1818 MacAlpine Road, R.R. #3, Gananoque, Ontario K7G 2V5 (CA). <b>(74) Agents:</b> STEEG, Carol, Miernicki et al.; Parteq Innovations, Queen's University at Kingston, Kingston, Ontario K7L 3N6 (CA).		<b>(81) Designated States:</b> AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>Without international search report and to be republished upon receipt of that report.</i>
<b>(54) Title:</b> METHODS FOR REMODELING NEURONAL AND CARDIOVASCULAR PATHWAYS		
<b>(57) Abstract</b> <p>The present invention provides a method of administration of an agent which acts to remodel neuronal or vascular pathways for the long term management of sexual dysfunction in both males and females. In a preferred embodiment, the invention provides a method of ameliorating or reversing pathogenic vascular degradative modeling in the ilio-hypogastric-pudental arterial bed and genitalia comprising administering to a human patient in need of such treatment a therapeutically effective amount of an anti-pressor agent. The anti-pressor agent comprises one or more compounds selected from the therapeutic classes of direct vasodilators such as hydralazine and NO donors, ACE inhibitors, angiotensin-II receptor antagonists, <math>\alpha_1</math>-adrenergic receptor antagonists, <math>\beta</math>-adrenergic receptor antagonists, calcium channel blockers, and phosphodiesterase inhibitors. The anti-pressor agent may be co-administered with a diuretic compound, and is administered either chronically at low dose, or for short periods of time at doses higher than are typically used for the treatment of hypertension. In certain embodiments of the method of the invention, the anti-pressor agent is co-administered with a diuretic agent and/or prostaglandin-E<sub>1</sub>.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon	KR	Republic of Korea	PL	Poland		
CN	China	KZ	Kazakstan	PT	Portugal		
CU	Cuba	LC	Saint Lucia	RO	Romania		
CZ	Czech Republic	LI	Liechtenstein	RU	Russian Federation		
DE	Germany	LK	Sri Lanka	SD	Sudan		
DK	Denmark	LR	Liberia	SE	Sweden		
EE	Estonia			SG	Singapore		

## Methods for Remodeling Neuronal and Cardiovascular Pathways

### Cross-Reference to Related Applications

5 This application claims priority to provisional application Serial No. 60/098,178 filed August 26, 1998.

### Technical Field

The present invention relates to medical methods of treatment, pharmaceutical compositions, and use of anti-pressor agents to manufacture such pharmaceutical  
10 compositions. More particularly, the present invention concerns the administration of an agent which acts to remodel neuronal or vascular pathways for the long term management of sexual dysfunction in both males and females.

### Background of the Invention

15 The physiology of an erection or sexual arousal in both the male and female involves central nervous system initiation, neural pathway activation, and vascular smooth muscle relaxation. This signaling mediates vasodilation of the penile, clitoral labial, and vaginal arterial blood vessels and the trabecular meshwork of smooth muscle. The resulting decrease in vascular resistance promotes an increase in arterial  
20 inflow and the filling of the *corpora cavernosa* in the penis and clitoris. Subsequent to there being an appropriate high rate of inflow, the cavernosal "filling" results in occlusion of the sub-tunical veins and full rigidity. The rate of inflow is critical because if there is not enough volume change, venous occlusion can not take place. A selective structurally-based increase in penile resistance produces a substantial impediment to  
25 inflow. That is, if penile or clitoral vascular structure, or the vascular structure immediately "up-stream" from the genitalia, is more constrained than the rest of the circulation, there would be a "mismatching" of perfusion pressure and selective resistance, i.e. genital arterial insufficiency. On the other hand, it is likely that when hypertension is first established and there is a generalized up-regulation of  
30 structurally-based vascular resistance in all vessels, there would not be any deleterious effect on erectile function because of a "matching" between perfusion pressure and

resistance. That is, despite the hypertrophy of the penile vasculature, the arterial pressure is proportionally elevated thereby allowing for adequate blood flow to the penis.

Pathological changes in the genital vasculature and alterations in function control systems have been shown to have a deleterious impact on erectile dysfunction. Local factors such as endothelin and sympathetic nerve mediated release of catecholamines have been shown to be important players in detumescence, but they also are likely to increase trophic responses in this tissue. The physiology of penile and clitoral erection and the structural maintenance of the tissue depends upon a balance between control systems that involve endothelial cells, vascular smooth muscle cells, fibroblasts, extracellular matrix, and nerves. Any shift in the balance of these control systems to either towards trophic responses such as vascular hypertrophy, focal fibrosis, or generalized production of the extracellular matrix or to the extremes of functional control systems can result in erectile dysfunction. Further, as structure and function are so closely related, it is becoming increasingly important in understanding the mechanisms of erectile dysfunction that we investigate the reciprocal impact of structural changes on function and of changes in functional control systems on structure.

The clitoris is the homologue of the penis, arising from the embryological genital tubercle. As a result, the two organs have similar structural and arousal response mechanisms. The clitoris consists of a cylindrical, erectile organ composed of three parts: the outermost glans or head, the middle corpus or body, and the innermost crura. The body of the clitoris consists of paired corpora cavernosa of about 2.5 cm in length and lacks a corpus spongiosum. During sexual arousal, blood flow to the corpora cavernosa of the clitoris cause their enlargement and tumescence.

The clitoris plays a major role during sexual activity in that it contributes to local autonomic and somatic changes causing vaginal vasocongestion, engorgement, and subsequent effects, lubricating the introital canal making the sexual act easier, more comfortable, and more pleasurable.

Vaginal wall engorgement enables a process of plasma transduction to occur, allowing a flow through the epithelium and onto the vaginal surface. Plasma

transduction results from the rising pressure in the vaginal capillary bed during the sexual arousal state. In addition, there is an increase in vaginal length and luminal diameter, especially in the distal 2/3 of the vaginal canal.

It has been well established that the generation of a penile and clitoral erections and vaginal and labial engorgement are greatly dependent on adequate blood flow to vascular beds which feed these organs. Both smooth muscle relaxation of the *corpora cavernosa* as well as the vasodilation of genital arterial vessels mediate the physiological response. One of the major fundamental etiologies of erectile dysfunction is, thus, inadequate genital arterial inflow. If there is an inappropriate structural narrowing in the supporting vasculature that is not associated with an increase in perfusion pressure, the blood flow into the organs at maximum dilation may be reduced and therefore be insufficient for the generation of an erection. There is increasing recognition that erectile dysfunction, although associated with, may appear prior to the onset of clinical signs of cardiovascular disease and therefore may be an early harbinger of progressing changes.

In both the male and female human, the aorta bifurcates on the fourth lumbar vertebra into the common iliac arteries. The common iliac arteries pass laterally, behind the common iliac veins, to the pelvic brim. At the lower border of the fifth lumbar vertebra, the common iliac arteries divide into internal and external branches. The internal iliac artery supplies blood to all of the organs within the pelvis and send branches through the greater sciatic notch to supply the gluteal muscles and perineum. After passing over the pelvic brim, the internal iliac artery divides into anterior and posterior trunks.

The anterior trunk of the internal iliac artery branches into the superior vesical artery, the inferior vesical artery, the middle rectal artery, the uterine artery, the obturator artery, the internal pudendal artery, and the inferior gluteal artery. The internal pudendal artery supplies blood to the perineum. The artery passes out of the pelvis around the spine of the ischium and back on the inside surface of the ischeal tuberosity and inferior ramus to lie in the pudendal canal. The branches from the internal pudendal artery are the inferior rectal artery which supplies the anal sphincter,

WO 00/12110

skin and lower rectum; the perineal artery which supplies the scrotum in the male and the labia in the female; the artery of the bulb which supplies erectile tissue, the deep dorsal arteries of the penis or deep artery of the clitoris.

It has been demonstrated in several forms of experimental hypertension that "slow pressor mechanisms" such as hypertrophic structural changes in the vasculature can almost completely account for the long-term resistance changes associated with the elevated arterial pressure. Based on Poiseuille's law, it has been shown that vascular resistance in an intact vascular bed is a function of the overall hemodynamic effect of all lumen radii, the number of blood vessels, the length of the vessels and the blood viscosity. In hypertension, increased vascular resistance is most potently conferred by a structurally-based decrease in the radius of the lumen of arterioles and small arteries and also potentially by arteriolar rarefaction whereby even a small change in the average arteriolar radii throughout a vascular bed has a dramatic influence on the resistance to flow. Further, it has been demonstrated that such structural changes can precede the onset of hypertension and therefore may be an initiating mechanism.

Vascular beds in which there is chronic diminished blood flow suffer a degree of pathogenic vascular degradative modeling over time in response to static or circulatory hypoxia. That is, as a normal reaction to diminished blood flow, the lumen in these arteries diminishes in diameter over time, causing decreased blood flow and/or higher pressure during periods of peak blood flow. Those portions of the ilio-hypogastric-pudendal arterial bed which directly feed blood to the sex organs are examples of such less frequently used arterial beds. Because incidents of major blood inflow to the sexual organs are less frequent than to most other organs, a gradual hypoxic response is seen over time in the vasculature directly feeding these organs and in the organs themselves. The body has a self-regulating mechanism to combat this pathogenic modeling: it is known, for example, that in the human male there are a number of spontaneous nocturnal erections which occur as a result of the body's mechanism for combating hypoxia in penile tissue. Nevertheless, the arteries in a normal flaccid penis and the un-enlarged clitoris and labia are constricted. As a result, typical oxygen concentrations in such tissues are closer to venous rather than arterial oxygen levels.



Periodic vasodilation of the penis and clitoris increases oxygen levels in these tissues. The higher oxygen levels supplied to tissue in the penis and clitoris, as well as vasodilation itself, shut down adverse metabolic processes such as TGF- $\beta$  production and pathogenic vascular wall modeling which result in long term tissue damage.

5           Therefore, it is differential changes in genital vascular resistance that is likely to be a critical issue in erectile function. That is, if such vascular structural changes take place in the genitalia in the absence of hypertension or systemic changes in vessel structure there would not be the increase in arterial pressure required to compensate for the increased resistance. It may be that this condition could occur as an early indicator  
10 of progressing cardiovascular disease. The appearance of erectile dysfunction preceding the global clinical signs of hypertension may, in fact, suggest an increased susceptibility of this vascular bed to pathological changes.

#### Summary of the Invention

15           In its principal embodiment, the present invention provides a method for the long term management of sexual dysfunction in males and females by administering a therapeutic agent which remodels neuronal or vascular pathways. In a preferred embodiment, the invention provides a method of ameliorating, inhibiting or reversing pathogenic vascular degradative modeling in the ilio-hypogastric-pudendal arterial bed  
20 and genitalia comprising administering to a human patient in need of such treatment a therapeutically effective amount of an anti-pressor agent. In one embodiment, the present invention provides the use of an anti-pressor agent for the manufacture of pharmaceutical compositions for ameliorating, inhibiting or reversing pathogenic vascular degradative modeling in the ilio-hypogastric-pudendal arterial bed and  
25 genitalia.

The anti-pressor agent is administered chronically at low doses ranging between about one-twentieth to about one-half the dose required to evoke vasodilation in a patient exhibiting normal circulation or, alternatively, is administered over a period of time ranging between about five days to about 21 days at higher doses ranging between

about 1.5 to about 3 times the dose required to evoke vasodilation in a patient exhibiting normal circulation.

Brief Description of the Drawing Figures

IN THE DRAWING:

- 5 **FIGURE 1** is a representative cumulative  $\alpha_1$ -adrenoreceptor concentration-response curve for administration of several doses of methoxamine (MXA) to a spontaneously hypertensive rat. Arrows indicate the point of drug delivery to the penile vascular bed at the concentrations labeled in the Figure. Each concentration of MXA was infused for a period of ten minutes, at which time a plateau was reached. The point marked "yield" in the Figure represents the pressure at maximum constriction of the blood vessels in the vascular bed. This maximum constriction was achieved by administration of a "cocktail" containing a mixture of vasopressin (20 g/mL), angiotensin-II (200 g/mL), and methoxamine (64 g/mL).
- 10
- 15 **FIGURE 2** shows the average  $\alpha_1$ -adrenoreceptor concentration-response curves for administration of methoxamine (MXA) to both the penile vascular bed and hindlimb vascular bed perfusion preparations of the spontaneous hypertensive rat (SHR) and the normotensive Sprague-Dawley rat (SD). Figure 2a and 2b represent, respectively, the curves for administration to the penile vascular beds of the SHR and SD rat strains. Figures 2c and 2d represent, respectively, the curves for administration to the hindlimb vascular beds of the SHR and SD rat strains.
- 20
- 25 **FIGURE 3a** shows the structurally-based vascular resistance asserted at maximum dilation for the penile and hindlimb perfusion vascular preparations for the spontaneously hypertensive rat (SHR) and the normotensive Sprague-Dawley rat (SD).
- 30 **FIGURE 3b** shows the corresponding structurally-based vascular resistance asserted at maximum constriction for the penile and hindlimb perfusion vascular preparations for the spontaneously hypertensive rat (SHR) and the normotensive Sprague-Dawley rat (SD).

FIGURE 4 is a schematic representation depicting structural differences in blood vessels in the spontaneously hypertensive rat (SHR) and the normotensive Sprague-Dawley rat (SD) and the expected impact on resistance to blood flow.

#### Detailed Description

5 The present invention contemplates the use of anti-pressor agents to remodel vasculature in the arterial beds supplying blood to the sex organs, and in the vascularity of the sex organs themselves. There has been some controversy in the literature as to the correct definition of the term "vascular remodeling," as evidenced by the exchange  
10 of letters in the Journal of Hypertension, 15: 333-337 (1997). The controversy in the nomenclature centers, in part, around the use of the terms "hypotrophic," "eutrophic," and "hypertrophic" as modifiers for the term "remodeling" as well as the use of the prefix "re-" in the word "remodeling."

15 The "trophic" terms have been objected to because of their suggestion that some sort of growth change accompanies the observed vascular changes. The term "remodeling" was initially applied in the literature to the observation in spontaneously hypertensive rats and in hypertensive humans that the interior lumen radius ( $r_i$ ) of blood vessels was greatly diminished while vessel wall mass ( $w$ ) remained constant. The  
20 result was an observed increase in the ratio of  $w/r_i$  which correlated with blood pressure elevation. The term "remodeling" was applied to the observed phenomenon, primarily because of the surprising consistency in total wall mass. It was thought that some sort of remodeling of the internal cellular structure of the blood vessel had occurred which permitted a change in lumen radius without a corresponding change in vessel wall mass.

25 The "re-" prefix has been objected to mainly because of the suggestion that some sort of "modeling" of the vasculature has already occurred, and subsequent changes (for good or ill) result in a "re-"modeling of those changes.

30 Lacking a general consensus of the term "vascular remodeling" in the medical community, the term "pathogenic vascular degradative modeling" will be applied, throughout this specification and the appended claims, to denote the pathogenic or degradative increase in the ratio  $w/r_i$  of vasculature, irrespective of the cause. The

term "vascular remodeling" as used throughout this specification and the appended claims will mean the amelioration, inhibition or reversal of pathogenic vascular degradative modeling; that is the amelioration, inhibition or reversal of the decrease in the ratio of vascular  $w/r_1$ .

5       The term "anti-pressor agent" as used herein denotes a therapeutic agent which acts either directly or indirectly to lower blood pressure. The term anti-pressor agent is chosen, rather than the more specific term "antihypertensive" agent, because the invention contemplates the use of agents which are effective to increase vascular flow in both hypertensive and normotensive patients. Anti-pressor agents contemplated for use  
10   in the method of the present invention include agents which act to bring about a lowering of blood pressure by any of a number of different physiological mechanisms. Anti-pressor agents include compounds belonging to a number of therapeutic classes based upon their mechanism of action, even though the therapeutic outcome is the same. Anti-pressor agents suitable for the method of this invention include compounds which  
15   are direct-acting vasodilators such as NO donors and hydralazine. Other suitable anti-pressor agents are compounds which act to inhibit the enzyme which converts the less potent decapeptide vasoconstrictor, angiotensin-I, to the more potent octapeptide vasoconstrictor, angiotensin II (so-called angiotensin-II converting enzyme inhibitors or "ACE inhibitors"), as well as agents which block the binding of angiotensin-II to the  
20   AT<sub>1</sub> receptor ("angiotensin-II receptor antagonists"). Anti-pressor agents useful in the method of the present invention also include vasodilating agents which act at  $\alpha_1$ -adrenergic receptors or  $\beta$ -adrenergic receptors in the smooth muscle of vascular walls (" $\alpha_1$ -adrenergic receptor antagonists" and " $\beta$ -adrenergic receptor antagonists"), as well as agents which act to decrease intracellular calcium ion concentration in arterial smooth  
25   muscle ("calcium channel blockers"). Suitable anti-pressor agents for use in the present invention also include activators of the enzymes guanylyl cyclase and adenylyl cyclase such as YC-1 and forskolin, respectively. PGE<sub>1</sub> (prostaglandin-E<sub>1</sub>), which acts both as an anti-pressor agent and as a sexual response initiator, is also suitable for use in the invention. Also contemplated as falling within the scope of the invention for use as  
30   anti-pressor agents are phosphodiesterase inhibiting agents, particularly type-3 and

type-5 phosphodiesterase inhibitors. Antagonists of PDE-5 (phosphodiesterase type 5), the enzyme primarily responsible for the degradation of cyclic guanosine monophosphate (cGMP), produce an increase in levels of cGMP, which, by way of "cross-talk," also decreases the activity of PDE-3, the enzyme primarily responsible for the degradation of cyclic adenosine monophosphate (cAMP). Thus, increasing levels of cGMP acts to inhibit the PDE-3 enzyme, thereby blocking the degradation of cAMP and causing an increase in cAMP levels. Thus, inhibition of either PDE-5 or PDE-3 results in an overall increase in concentrations of cAMP and cGMP.

Specific examples of NO donors include glyceryl trinitrate, isosorbide 5-mononitrate, isosorbide dinitrate, pentaerythritol tetranitrate, sodium nitroprusside, 3-morpholiniosydnonimine, molsidnomine, S-nitroso-N-acetylpenicillamine, S-nitrosoglutathione, N-hydroxyl-L-arginine, S,S-dinitrosodithiol, and NO gas.

ACE inhibitors include benzazapine compounds such as benazepril, and libenzapril; 6H-pyridazino[1,2-a]diazepine derivatives such as cilazapril; 2,3-dihydro-1H-indene compounds such as delapril; L-proline derivatives such as alacepril, captopril, ceronapril, enalapril, fosinopril, lisinopril, moveltipril and spirapril; oxoimidazoline derivatives such as imidapril; 1,4-dihydropyridine compounds such as lacidipine; *iso*-quinoline carboxylic acid derivatives such as moexipril and quinapril; 1H-indole carboxylic acid derivatives such as pentopril and perindopril; hexahydroindole carboxylic acid derivatives such as trandolapril; cyclopenta[b]pyrrole carboxylic acid derivatives such as ramipril; and 1,4-thiazepine compounds such as temocapril.

Angiotensin-II receptor antagonists useful as anti-pressor agents in the method of this invention include eprosartan, irbesartan, losartan, and valsartan.

$\alpha_1$ -Adrenergic receptor antagonists include substituted phenyl derivatives such as midrodrine, phenoxybenzamine, tamsulosin; substituted naphthyl derivatives such as naphazoline; aminoquinazoline derivatives such as alfuzosin, bunazosin, doxazosin, prazosin, terazosin and trimazosin; benzamide compounds such as labetolol; carbazole derivatives such as carvedilol; dimethyluracil derivatives such as urapidil; imidazolidinyl derivatives such as apraclonidine, clonidine; dihydroimidazole

derivatives such as phentolamine; indole derivatives such as indoramin; and 1,2,4-triazolo[4,3- $\alpha$ ]pyridine compounds such as dapiprazole.

Calcium channel blockers include benzothiazepine compounds such as diltiazem; dihydropyridine compounds such as nifedipine, nifedipine, and nimopidine; 5 phenylalkylamine compounds such as verapamil; diarylaminopropylamine ether compounds such as bepridil; and benimidazole-substituted tetralin compounds such as nifedipine.

Phosphodiesterase inhibitors include bipyridone compounds such as amrinone; and dihydropyrazolopyrimidine compounds such as sildenafil. Sildenafil functions as a 10 selective type-5 (i.e. c-GMP specific) phosphodiesterase inhibitor, and acts to decrease the metabolism of c-GMP, the second messenger in nitric oxide mediated erectile response. An oral formulation of this medication has proven to be safe and effective in improving erectile duration and rigidity. In females, nitric oxide/NOS exists in human vaginal and clitoral tissue. Sildenafil, alone or in combination with other vasoactive 15 agents, is effective for the long term management of sexual dysfunction for the treatment of vasculogenic male or female sexual dysfunction.

#### Pharmaceutical Compositions

Pharmaceutical compositions which are useful in the method of the present 20 invention comprise one or more compounds defined above formulated together with one or more non-toxic pharmaceutically acceptable carriers. The pharmaceutical compositions may be specially formulated for oral administration in solid or liquid form, for parenteral injection, or for vaginal or rectal administration. The formulations may, for example, contain a single therapeutic agent selected from ACE inhibitors, 25 angiotensin-1 ( $AT_1$ ) receptor antagonists,  $\alpha_1$ -adrenoreceptor antagonists,  $\beta$ -adrenergic receptor antagonists, direct-acting vasodilators, NO donors, calcium channel blockers, phosphodiesterase inhibitors, or a combination of two or more agents selected from the same or different therapeutic categories. Moreover, a combination of one or more therapeutic agents from the groups listed above may be combined with a diuretic agent 30 of the class well known in the art.

To enhance delivery to genital vasculature, combined systemic delivery with topical administration of an erectogenic initiator is also contemplated within the scope of this invention. In this manner the anti-pressor drug is delivered to target regions at a markedly enhanced rate. Since prostaglandin- $E_1$  acts both as an anti-pressor and as a direct sexual response initiator, one or more therapeutic agents from the groups listed above can be administered in combination therapy with prostaglandin  $PGE_1$ . The co-administered  $PGE_1$  may be administered by any of the routes discussed below, with topical application being a preferred route.

The pharmaceutical compositions of this invention can be administered to either systemically or locally to humans and other animals. Systemic routes include oral, parenteral, intracisternal, intraperitoneal, trans-cutaneous (by injection or patch), buccal, sub-lingual administration, or by means of an oral or nasal spray. The term "parenteral" administration as used herein refers to modes of administration which include intravenous, intramuscular, intraperitoneal, intrasternal, subcutaneous and intraarterial injection and infusion. Local administration routes include vaginal, rectal, intraurethral, trans-urethral, by intra-cavernosal injection, or topical administration.

Pharmaceutical compositions of this invention for parenteral injection comprise pharmaceutically acceptable sterile aqueous or nonaqueous solutions, dispersions, suspensions or emulsions as well as sterile powders for reconstitution into sterile injectable solutions or dispersions just prior to use. Examples of suitable aqueous and nonaqueous carriers, diluents, solvents or vehicles include water, ethanol, polyols (such as glycerol, propylene glycol, polyethylene glycol, and the like), and suitable mixtures thereof, vegetable oils (such as olive oil), and injectable organic esters such as ethyl oleate. Proper fluidity can be maintained, for example, by the use of coating materials such as lecithin, by the maintenance of the required particle size in the case of dispersions, and by the use of surfactants.

These compositions may also contain adjuvants such as preservatives, wetting agents, emulsifying agents, and dispersing agents. Prevention of the action of microorganisms may be ensured by the inclusion of various and bacterial and antifungal agents, for example, paraben, chlorobutanol, phenol sorbic acid, and the like. It may

also be desirable to include isotonic agents such as sugars, sodium chloride, and the like. Prolonged absorption of the injectable pharmaceutical form may be brought about by the inclusion of agents which delay absorption such as aluminum monostearate and gelatin.

5 In some cases, in order to prolong the effect of the drug it is desirable to slow the release or absorption of the drug following subcutaneous or intramuscular injection. This may be accomplished by the use of a liquid suspension of crystalline or amorphous material with low water solubility. The rate of absorption of the drug then depends upon its rate of dissolution which, in turn, may depend upon crystal size and crystalline form.  
10 Alternatively, delayed absorption of a parenterally administered drug form is accomplished by dissolving or suspending the drug in an oil vehicle.

Injectable depot forms are made by forming microencapsule matrices of the drug in biodegradable polymers such as polylactide-polyglycolide. Depending upon the ratio of drug to polymer and the nature of the particular polymer employed, the rate of drug  
15 release can be controlled. Examples of other biodegradable polymers include poly(orthoesters) and poly(anhydrides). Depot injectable formulations are also prepared by entrapping the drug in liposomes or microemulsions which are compatible with body tissues.

The injectable formulations can be sterilized, for example, by filtration through  
20 a bacterial-retaining filter, or by incorporating sterilizing agents in the form of sterile solid compositions which can be dissolved or dispersed in sterile water or other sterile injectable medium just prior to use.

Solid dosage forms for oral administration include capsules, tablets, pills, powders, and granules. In such solid dosage forms the active compound is mixed with  
25 at least one inert, pharmaceutically acceptable excipient or carrier such as sodium citrate or dicalcium phosphate and/or a) fillers or extenders such as starches, lactose, sucrose, glucose, mannitol, and silicic acid, b) binders such as, for example, carboxymethylcellulose, alginates, gelatin, polyvinylpyrrolidone, sucrose, and acacia, c) humectants such as glycerol, d) disintegrating agents such as agar-agar, calcium  
30 carbonate, potato or tapioca starch, alginic acid, certain silicates, and sodium carbonate, e) solution retarding agents such as paraffin, f) absorption accelerators such as



quaternary ammonium compounds, g) wetting agents such as, for example, cetyl alcohol and glycerol monostearate, h) absorbents such as kaolin and bentonite clay, and l) lubricants such as talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate, and mixtures thereof. In the case, of capsules, tablets and pills, the dosage form may also comprise buffering agents. Solid compositions of a similar type may also be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugar as well as high molecular weight polyethylene glycols and the like.

The solid dosage forms of tablets, dragees, capsules, pills, and granules can be prepared with coatings and shells such as enteric coatings and other coatings well known in the pharmaceutical formulating art. They may optionally contain opacifying agents and can also be of a composition that they release the active ingredient(s) only, or preferentially, in a certain part of the intestinal tract, optionally, in a delayed manner. Examples of embedding compositions which can be used include polymeric substances and waxes. The active compounds can also be in micro-encapsulated form, if appropriate, with one or more of the above-mentioned excipients.

Liquid dosage forms for oral administration include pharmaceutically acceptable emulsions, solutions, suspensions, syrups and elixirs. In addition to the active compounds, the liquid dosage forms may contain inert diluents commonly used in the art such as, for example, water or other solvents, solubilizing agents and emulsifiers such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, dimethyl formamide, oils (in particular, cottonseed, ground nut corn, germ olive, castor, and sesame oils), glycerol, tetrahydrofurfuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan, and mixtures thereof. Besides inert diluents, the oral compositions can also include adjuvants such as wetting agents, emulsifying and suspending agents, sweetening, flavoring, and perfuming agents.

Suspensions, in addition to the active compounds, may contain suspending agents as, for example, ethoxylated isostearyl alcohols, polyoxyethylene sorbitol and

sorbitan esters, microcrystalline cellulose, aluminum metahydroxide, bentonite, agar-agar, and tragacanth, and mixtures thereof.

Compositions for rectal administration are preferably suppositories which can be prepared by mixing the compounds of this invention with suitable non-irritating excipients or carriers such as cocoa butter, polyethylene glycol or a suppository wax which are solid at room temperature but liquid at body temperature and therefore melt in the rectum and release the active compound,

Compounds of the present invention can also be administered in the form of liposomes. As is known in the art, liposomes are generally derived from phospholipids or other lipid substances. Liposomes are formed by mono- or multi-lamellar hydrated liquid crystals that are dispersed in an aqueous medium. Any non-toxic, physiologically acceptable and metabolizable lipid capable of forming liposomes can be used. The present compositions in liposome form can contain, in addition to a compound of the present invention, stabilizers, preservatives, excipients, and the like. The preferred lipids are the phospholipids and the phosphatidyl cholines (lecithins), both natural and synthetic. Methods for the formation of liposomes are known in the art. See, for example, Prescott, Ed., Methods in Cell Biology, Volume XIV Academic Press, New York, N.Y. (1976), p. 33 et seq.

Actual dosage levels of active ingredients in the pharmaceutical compositions of this invention may be varied so as to obtain an amount of the active compound(s) that is effective to achieve the desired therapeutic response for a particular patient, compositions, and mode of administration. The selected dosage level will depend upon the activity of the particular compound, the route of administration, the severity of the condition being treated, and the condition and prior medical history of the patient being treated. However, it is well known within the medical art to determine the proper dose for a particular patient by the "dose titration" method. In this method, the patient is started with a dose of the drug compound at a level lower than that required to achieve the desired therapeutic effect. The dose is then gradually increased until the desired effect is achieved. Starting dosage levels for an already commercially available therapeutic agent of the classes discussed above can be derived from the information

already available on the dosages employed for the use of the compound as an antihypertensive agent. In a chronic, or long-term dosing regimen to remodel the vasculature in the genitalia and in vascular beds feeding the genitalia, lower doses ranging between about 1/20 to about 1/2 the doses normally given to combat hypertension are used. In short term, acute, or "burst-mode" therapy, the compounds are administered in doses ranging between 1 to 3 times the amounts generally prescribed for hypertension. In these situations, however, appropriate precautions should be taken by the attending physician to closely monitor untoward side-effects peculiar to each particular therapeutic agent.

For the preferred therapeutic agents in the method of the present invention, namely ACE inhibitors, generally dosage levels of about 1 mg to about 300 mg, more preferably of about 5 mg to about 150 mg of active compound per kilogram of body weight per day are administered orally to a patient, with the dose levels appropriately adjusted if the route of administration is other than oral. If desired, the effective daily dose may be divided into multiple doses for purposes of administration, e.g. two to four separate doses per day.

#### Biological Data

##### A. Demonstration that the Sex Organs are Not Protected from Pathological Vascular Degradative Modeling

###### 1 Methodology

Male spontaneously hypertensive (SH) rats weighing between 246-313 g, and normotensive Sprague-Dawley (SD) rats weighing between 246-440 g were obtained from Charles River Laboratories (Montreal, Quebec, Canada). The animals were maintained in individual cages with a 12 hour light / 12 hour dark cycle, and a room temperature of 22-24°C. They were provided with standard rodent chow and tap water *ad libitum* and were acclimated to the room for at least two days before the experiments. All procedures were carried out in accordance with the guidelines set out by the Canadian Council on Animal Care.

## 2. Penile vascular resistance properties

Penile perfusion preparations were made using the procedure established by Banting, J. D., *et al.*, "Isolation and Perfusion of the Pudendal Vasculature in Male Rats. *J Urol.*, 2: 587-590 (1995). A heated chamber served to maintain the ambient temperature and the entire preparation at 37-38°C. The perfusate was held in a reservoir and passed through a heating and a bubble trapping/mixing chamber connected to a single peristaltic pump (Minipuls 2 Pump, Gilson, Inc., 3000 W. Beltline Highway, Middleton, WI USA 53562). An injection port was located distal to the pump for the introduction of pharmacological agents to minimize dead volume. Drugs were administered via a Harvard Apparatus, Inc. infusion pump (Harvard Apparatus, Inc., 84 October Hill Road, Holliston, Massachusetts 01746). The perfusate was a Tyrode-dextran solution consisting of a mixture of 20 mg of KCl, 32.3mg of  $\text{CaCl}_2 \cdot \text{H}_2\text{O}$ , 5.1 mg of  $\text{MgCl}_2 \cdot 6\text{H}_2\text{O}$ , 6.2 mg of  $\text{NaH}_2\text{PO}_4 \cdot 2\text{H}_2\text{O}$ , 155 mg of  $\text{NaHCO}_3$ , 100 mg of glucose, and 800 mg of NaCl in each 100 mL of fluid. The solution was maintained at pH 7.4, and a temperature of 37-39°C, and oxygenated with 95%  $\text{O}_2$  and 5%  $\text{CO}_2$ . The rats were anaesthetized (sodium pentobarbital 60 mg/kg body weight i.p.) and heparinized (1000 IU/kg, i.v.). The bilateral isolation of penile vasculature was achieved by ligating all of the branching arteries except for the pudendal; then the abdominal aorta was cannulated proximal to the iliac bifurcation with a single lumen catheter. The catheter was connected to the perfusion apparatus via a pressure transducer for arterial pressure recording. After sectioning the *vena cava* and spinal cord to remove venous resistance and to eliminate neural influences, a flow of perfusate (1 mL/min per kg body weight) through the abdominal cannula was initiated. The perfusion pressure was continuously recorded on a data acquisition system (MacLab, AD Instruments, Houston, TX). The perfusate was infused for twenty minutes to flush the penile vasculature of blood and obtain a stable pressure before the beginning of any experiment. Following this, sodium nitroprusside (20 g/mL) was infused to induce maximum vasodilation. The flow rate-perfusion/pressure relationship was determined by measuring the pressure at minimum vascular resistance at flow rates of 0.5, 1.0, 2.0, 4.0 mL/min per kg of body weight. A cumulative  $\alpha_1$ -adrenoreceptor

concentration-response curve to methoxamine (2.5, 5, 10, 25, 50 g/mL was then generated. Each concentration of methoxamine was infused for a duration of 10 minutes, at which time a plateau was reached. Subsequently, a continuous injection of a cocktail containing a supramaximal concentration of vasoconstrictors (vasopressin, 20.5 g/mL, angiotensin-II, 200 ng/mL; methoxamine, 64 g/mL; Sigma, St. Louis, MO, 63178) was given to ensure that maximum constrictor response that was not dependent upon the activation of a single receptor type was achieved. A second injection of the constrictor cocktail was administered to ensure the tissue "yield" was maximum constriction. This "yield" induced by the multi-vasoconstrictor cocktail has been demonstrated to correlate directly with the bulk of medial vascular smooth muscle cells in the resistance vasculature. A typical perfusion pressure tracing from this protocol can be seen in Figure 1. At the end of the concentration-response relationship, the aorta was cut distal to the catheter, and a baseline flow-pressure curve was recorded again. This was done to ensure that pressure fell to zero and to account for any false pressure readings that may have resulted due to movement of the catheter during the experiment.

### 3. Hindlimb vascular resistance properties

The hindlimb perfusion preparation was adopted from a technique originally designed by Folkow *et al.*, Acta Physiol Scand., 80: 93-106 (1973), as modified by Adams *et al.*, Hypertension, 14: 191-202 (1989). The perfusion experiment was performed as described above. Drugs were administered into the mixing chamber via a Harvard Apparatus infusion pump. The rats were anaesthetized (Inactin, 100 mg/kg of body weight, i.p.) and heparinized (1000 IU/kg of body weight, i.v.). Following a midline abdominal incision, the abdominal aorta was cannulated proximal to the iliac bifurcation with a double lumen catheter (Storz, St Louis, MO, USA), and the catheter was extended down the right common iliac artery. One lumen of the catheter was connected to the perfusion apparatus, while the other was connected to a pressure transducer for arterial pressure recording. The rat was perfused at a constant flow rate (2 mL/min per 100g of body weight) and the experiments were carried out as described above. The flow rate/perfusion pressure relationship was recorded at flow rates of 0.5, 1.0, 2.0,

4.0 mL/min per 100 g of body weight. A cumulative  $\alpha_1$ -adrenoreceptor concentration-response curve to methoxamine (0.5, 1, 2, 4, 8, 16, 32, 64 g/mL) was then generated. Each concentration of methoxamine was infused for a duration of 5 minutes, at which time a plateau was reached. Subsequently, a bolus injection of a cocktail containing a supramaximal concentration of vasoconstrictors was given as above. At the end of the concentration-response relationship, the iliac artery was cut distal to the catheter, and the flow pressure curve was monitored again.

Flow rates for the hindlimb perfusion experiments were determined based on expected flow rates of exercising skeletal muscle at maximum dilation. The flow rate used resulted in a perfusion pressure at maximum dilation between 20-25 mm Hg which is well within the expected range. After checking several flow rates in the penile perfusion, a rate was obtained that resulted in a similar perfusion pressure at maximum dilation. The flow rates chosen also allow for the assessment at maximum constriction. This allowed for comparison between strains.

#### 4. Analysis of Data

All values in the figures and tables were expressed as mean  $\pm$  standard deviation. Results comparing penile and hindlimb vasculature were analyzed using the Student's t-test. Differences were considered as significant at  $p < 0.05$ .

#### 5. Results

There was no significant difference in the body weight of the spontaneously hypertensive rats in the penile assessment group ( $267 \pm 29$  g,  $n=5$ ) and in the hindlimb assessment group ( $270 \pm 5.7$  g,  $n=3$ ). The average body weight of the normotensive Sprague-Dawley rats in the hindlimb assessment group ( $375 \pm 41$  g,  $n=8$ ) was significantly higher than that of the of the normotensive Sprague-Dawley rats in the penile assessment group ( $284 \pm 32$  g,  $n=5$ ). However, this was not considered relevant because in normotensive adult rats there has been shown to be very little correlation between body weight and blood pressure (Adams, M. A., *et al.*, Hypertension, 14: 191-202 (1989)).

This hemodynamic analysis had similar effects in most parameters between the penile and hindlimb vascular beds within each rat strain. The flow-pressure curve assessed at maximal dilation was similar in both the penile and the hindlimb vasculatures of spontaneously hypertensive and normotensive rats as shown in Table 1.

- 5 These curves were monitored to ensure a linear increase in perfusion pressure with an increase in flow rate. The increase in the flow rate exerted a radial pressure against the vessel wall and resulted in increased pressure. The spontaneously hypertensive rats trended towards a higher baseline pressure than the normotensive rats. This was observed in both penile and hindlimb vascular beds. These data suggest that
- 10 spontaneously hypertensive rats may have a smaller lumen thus causing them to operate at a higher pressure than normotensive Sprague-Dawley rats even when there is no constrictor tone on the vessel.

Table 1

Group	Slope Flow Pressure	Maximum Constriction With Methoxamine (mm Hg)	Log EC <sub>50</sub>	Slope Methoxamine
SHR, penile bed	7.15 ± 2.0	172 ± 32	0.95 ± 0.19	1.64 ± 0.21
SHR, hindlimb bed	6.68 ± 0.38	253 ± 25*	0.79 ± 0.15	5.19 ± 3.0*
SD, penile bed	7.34 ± 2.3	171 ± 36	0.63 ± 0.24	2.03 ± 0.68
SD, hindlimb bed	6.99 ± 3.4	191 ± 55	0.78 ± 0.12	3.0 ± 0.99

\* Statistically significant.

- 15 Table I shows there was a statistically significant difference in maximum constriction with a supramaximal dose of methoxamine (50 g/mL for penile and 64 g/mL for hindlimb vasculature) between spontaneously hypertensive rat hindlimb vasculature (253 ± 25 mmHg) and spontaneously hypertensive rat penile vasculature (172 ± 32 mmHg). This difference was not observed in normotensive Sprague-Dawley
- 20 rats. The discrepancy is novel and requires further assessment. It is expected that higher responses would be seen in the spontaneously hypertensive rats in both arterial beds, however only the hindlimb vasculature showed a significant difference between

spontaneously hypertensive and normotensive rats. Average concentration response curves for methoxamine of the two strains in both beds are shown in Figures 2a-2d.

The  $EC_{50}$  of the concentration response curve shown in Table 1 gives the concentration of drug at which there is a 50% response to  $\alpha_1$ -adrenoreceptor stimulation. This value would be an indication of the sensitivity of the tissue to this receptor activation. The logs  $EC_{50}$  of the methoxamine concentration-response curves were not different for penile and hindlimb vasculature in both the spontaneously hypertensive and normotensive rats thus indicating similar sensitivity to this receptor stimulation.

The steepest slope of this curve is given in Table 1. In normotensive rats, there was no statistically significant difference in slope between penile vasculature ( $2.03 \pm 0.68$ ) and hindlimb vasculature ( $3.0 \pm 0.99$ ). The parameters showed a statistically significant difference between spontaneously hypertensive rat penile ( $1.64 \pm 0.21$ ) and spontaneously hypertensive rat hindlimb ( $5.19 \pm 3.0$ ). This was expected since the maximal constriction with methoxamine was lower in penile vasculature while the  $EC_{50}$  remained the same.

Figure 3 depicts the structurally-based vascular resistance properties assessed at both maximum dilation and maximum constriction. There was no significant difference in perfusion pressures at maximum dilation within the rat strains. Between the two strains of rats, the penile vasculature trended towards higher pressures in the hypertensive rat as compared to the normotensive Sprague-Dawley rats, however it did not reach a level of significance as in the hindlimb. Spontaneously hypertensive rats reached a point of maximal constriction with a cocktail at a perfusion pressure that was 20% higher than normotensive rats in each vascular bed. There was no statistically significant difference between the two beds within strain, suggesting that the penile and hindlimb vasculature undergo similar structural changes in genetically hypertensive rats. This point demonstrates the increased medial thickening that occurs in the hypertensive rats that allows for the maintenance of higher arterial operating pressures.



## 6. Discussion

The major findings of the data presented above demonstrate that the penile vasculature is not protected from the structural changes that take place in the other vascular beds of spontaneously hypertensive rats relative to normotensive strains.

5 Increased medial thickening and narrowing of the vascular lumen have been found in blood vessels of a wide range of vascular beds of spontaneously hypertensive rats. Therefore, the overall results of the present series of experiments have shown that the genetic disposition appears to dominate the structure regardless of the vascular bed.

In the present study a hemodynamic methodology was used to compare and  
10 contrast structurally-based vascular resistance in two vascular beds. The hindlimb bed was chosen for comparison since the vascular resistance properties are well established and anatomically the feeder vessels of the two beds are common.

These results demonstrate that the resistance properties at maximum dilation were similar in the two beds within strains. A general finding of studies comparing  
15 vascular resistance at minimum tone is that a higher perfusion pressure is normally obtained in SHR compared to normotensive rats. Thus, findings of elevated resistance properties at maximum dilation are consistent with there being an overall narrowing of the vascular lumen. Resistance properties were further assessed by determining the slope of the flow-pressure curve at maximum dilation. This relationship was used to  
20 determine whether there were any differences in the passive vascular wall elements such as the extracellular matrix components, i.e. if distensibility was altered there would be a differential effect on the flow-pressure curves. Further, a thicker medial wall could also result in a stiffer vessel which would exhibit less compliance with increasing flow. The lack of difference in all of these values suggests that there has been no differential  
25 change in the components of the vessel wall within the penile vasculature.

Assessment of the active components of the vessel walls was determined by inducing a state of maximal vasoconstrictor tone using a cocktail of vasoconstrictor agonists. The supramaximal, multiple agonist stimulus produces a maximum constrictor response which is independent of individual receptor population changes  
30 thereby reflecting only the overall contractile bulk of the medial smooth muscle cells.

The findings that sensitivity ( $EC_{50}$ ) and reactivity (slope) to  $\alpha_1$ -adrenoreceptor stimulation were not different between vascular beds or strains likely indicates that there is a similar stimulus-response coupling of the noradrenergic innervation in all of these vessels; i.e. there is a consensus of normal vascular biology. In the schematic diagram of Figure 4, the concept of structural changes dominating function is depicted. Thus, the curves show that, at any given level of vasoconstrictor tone, the hypertensive circulation will always have increased vascular resistance compared to normotensive circulation. Another way of looking at this concept would be that at the same level of constrictor tone, the normotensive circulation would be able to achieve the same inflow at a proportionately lower perfusion pressure than the hypertensive circulation.

Taken together, the findings indicate that the penile vasculature has an increased average medial mass coupled with decreased average lumen. The generation of an erection is based on the flow when vessels are in a dilated state. Although there was a significant difference in the perfusion pressure at maximum dilation in the hindlimb vasculature of spontaneously hypertensive rats (SHR) when compared to the normotensive Sprague-Dawley (SD) rats, this was not detected in the penile vascular bed. There was however a trend toward significance which may be seen in future studies when animals are used that are genetically closer to the spontaneously hypertensive rat, such as the Wistar-Kyoto rat (Taconic, 273 Hover Avenue, Germantown, NY 12526) which is a more appropriate normotensive control. The penile vasculature is more complex than that of the hindlimb and therefore it may be that differences at maximum dilation are more difficult to detect. It is also unknown whether the size of the penis differs between the strains examined in this study. The length of the vessels changes baseline resistance more than the maximum constriction response because the maximum dilation is flow-dependent, as there is no constrictor tone on the vessel. In contrast, maximum constriction responses are based on the bulk muscular tissue. Therefore although the point of maximum dilation is expected to be higher in the penile vasculature of SHR as compared to a normotensive control it may not be detectable using the Sprague-Dawley rats as a comparison based on the suspected differences between the strains.

## **B. Therapeutically-Induced Vascular Remodeling in Penile Vasculature**

### **1. Methodology**

Adult spontaneously hypertensive rats (SHR) were treated for 1 or two weeks with either enalapril (30 mg/kg of body weight per day) or hydralazine (45 mg/kg of body weight per day). Following this treatment, structurally-based resistance to blood flow in the perfused penile vascular bed and hindlimb vascular bed models were measured using the methods detailed above. Cumulative  $\alpha_1$ -adrenoreceptor concentration-response curves in response to methoxamine were measured as described above, and the "yield" point determined, following the achievement of maximal vasoconstriction using the vasopressin, angiotensin-II, methoxamine cocktail described above. The animal hearts were excised and weighed. The data are presented in Table 2 below.

**Table 2**

<b>Group</b>	<b>Slope Flow Pressure</b>	<b>Maximum Constriction With Cocktail "Yield" (mm Hg)</b>	<b>Log EC<sub>50</sub></b>	<b>Left Ventricle Weight (g)/ Body Weight (kg) Ratio</b>
SHR-E <sub>2</sub> <sup>1</sup> (n = 7)	6.45 ± 1.31	335 ± 23	8.73 ± 0.26	2.13 ± 0.08
SHR-E <sub>1</sub> <sup>1</sup> (n = 5)	6.10 ± 1.5	358 ± 20	7.33 ± 1.39	2.17 ± 0.05
SHR-H <sup>2</sup> (n = 7)	6.63 ± 0.86	353 ± 11	13.56 ± 5	2.37 ± 0.12
Control (n = 9)	7.13 ± 0.63	381 ± 21	11.95 ± 5.51	2.46 ± 0.08

<sup>1</sup> Enalapril-treated animals.

<sup>2</sup> Hydralazine-treated animals

The data appearing in Table 2 show that enalapril treatment progressively regressed (remodeled) cardiac and pudendal vascular structure over the 2-week period of treatment. The "yield" value decreased on average by 12.1% ± 6.0%, while left

ventricular mass decreased by  $13.6\% \pm 3.2\%$ . Hydralazine treatment was somewhat less effective, decreasing the "yield" point by  $7.3\% \pm 2.9\%$ , and had no significant effect on left ventricular weight (decreased of  $3.7\% \pm 5.0\%$ ).

While there have been shown and described what are believed at present to be the preferred embodiments of the present invention, it will be obvious to those of ordinary skill in the art that various modifications can be made in the preferred embodiments without departing from the scope of the invention as it is defined by the appended claims.

**WE CLAIM:**

1. A method of ameliorating, inhibiting or reversing pathogenic vascular degradative modeling in the ilio-hypogastric-pudendal arterial bed and genitalia  
5 comprising administering to a human patient in need of such treatment a therapeutically effective amount of an anti-pressor agent.
2. The method according to Claim 1 wherein said anti-pressor agent comprises one or more compounds selected independently from the group consisting of prostaglandin-  
10 E<sub>1</sub>, ACE inhibitors, AT<sub>1</sub>-receptor antagonists,  $\alpha_1$ -adrenergic receptor antagonists,  $\beta$ -adrenergic receptor antagonists, calcium channel blockers, direct acting vasodilators, NO donors, activators of guanylyl cyclase, activators of adenylyl cyclase, and phosphodiesterase inhibitors.
- 15 3. The method of Claim 2 wherein said anti-pressor agent is an activator of guanylyl cyclase or adenylyl cyclase selected from the group consisting of YC-1 and forskolin.
4. The method of Claim 2 wherein said anti-pressor agent is a direct acting  
20 vasodilator selected from the group consisting of hydralazine and NO donors.
5. The method of Claim 4 wherein said NO donor is selected from the group consisting of glyceryl trinitrate, isosorbide 5-mononitrate, isosorbide dinitrate, pentaerythritol tetranitrate, sodium nitroprusside, 3-morpholinodnonimine,  
25 molsidnomine, S-nitroso-N-acetylpenicillamine, S-nitrosoglutathione, N-hydroxyl-L-arginine, S,S-dinitrosodthiol, and NO gas.
6. The method according to Claim 2 wherein said anti-pressor agent is an angiotensin-II receptor antagonists selected from the group consisting of eprosartan,  
30 irbesartan, losartan, and valsartan, and mixtures thereof.

7. The method according to Claim 2 wherein said anti-pressor agent is an ACE inhibitor selected from the group consisting of alacepril, benazepril, captopril, ceronapril, cilazapril, delapril, enalapril, fosinopril, imidapril, lacidipine, libenzapril, lisinopril, moexipril, moveltipril, pentopril, perindopril, quinapril, ramipril, spirapril, temocapril, and trandolapril, and mixtures thereof.
8. The method according to Claim 2 wherein said anti-pressor agent is an  $\alpha_1$ -adrenergic antagonist selected from the group consisting of alfuzosin, apraclonidine, bunazosin, carvedilol, clonidine, dapiprazole, doxazosin, indoramin, labetolol, midrodrine, naphazoline, phenoxybenzamine, phentolamine, prazosin, tamsulosin, terazosin, trimazosin, and urapidil, and mixtures thereof.
9. The method according to Claim 2 wherein said anti-pressor agent is a calcium channel blocker selected from the group consisting of bepridil, diltiazem, mibrefadil, nicardipine, nifedipine, nimopidine, and verapamil, and mixtures thereof.
10. The method according to Claim 2 wherein said anti-pressor agent is a phosphodiesterase inhibitor selected from the group consisting of amrinone and sildenafil.
11. The method according to Claim 2 wherein said anti-pressor agent is co-administered with a diuretic compound.
12. The method according to Claim 2 wherein said anti-pressor agent is co-administered with prostaglandin- $E_1$ .
13. The method of Claim 1 wherein said anti-pressor agent is administered to a normotensive patient.

14. The method of Claim 1 wherein said anti-pressor agent is administered on a chronic basis at a dose ranging between one-twentieth to one-half the dose normally given to a hypertensive patient.
- 5 15. The method of Claim 1 wherein said anti-pressor agent is administered for a period ranging between about three days to about twenty-one days at a dose ranging between one to three times the dose normally given to a hypertensive patient.
16. The method according to Claim 14 wherein said anti-pressor agent is  
10 administered to a normotensive patient.
17. The method according to Claim 15 wherein said anti-pressor agent is administered to a normotensive patient.
- 15 18. A method of remodeling the vascular bed which supplies blood to the genitalia, comprising administering to a patient in need thereof a therapeutically effective amount of an anti-pressor agent.
19. The method according to Claim 18 wherein said anti-pressor agent comprises  
20 one or more compounds selected independently from the group consisting of prostaglandin-E<sub>1</sub>, ACE inhibitors, AT<sub>1</sub>-receptor antagonists,  $\alpha_1$ -adrenergic receptor antagonists,  $\beta$ -adrenergic receptor antagonists, calcium channel blockers, direct acting vasodilators, NO donors, activators of guanylyl cyclase, activators of adenylyl cyclase, and phosphodiesterase inhibitors.
- 25 20. The method of Claim 19 wherein said anti-pressor agent is an activator of guanylyl cyclase or adenylyl cyclase selected from the group consisting of YC-1 and forskolin.

21. The method of Claim 19 wherein said anti-pressor agent is a direct acting vasodilator selected from the group consisting of hydralazine and NO donors.
22. The method of Claim 21 wherein said NO donor is selected from the group consisting of glyceryl trinitrate, isosorbide 5-mononitrate, isosorbide dinitrate, pentaerythritol tetranitrate, sodium nitroprusside, 3-morpholinodinitrate, molsidomine, S-nitroso-N-acetylpencillamine, S-nitrosoglutathione, N-hydroxyl-L-arginine, S,S-dinitrosodithiol, and NO gas.
23. The method according to Claim 19 wherein said anti-pressor agent is an angiotensin-II receptor antagonists selected from the group consisting of eprosartan, irbesartan, losartan, and valsartan, and mixtures thereof.
24. The method according to Claim 19 wherein said anti-pressor agent is an ACE inhibitor selected from the group consisting of alacepril, benazepril, captopril, ceronapril, cilazapril, delapril, enalapril, fosinopril, imidapril, lacidipine, libenzapril, lisinopril, moexipril, moveltipril, pentopril, perindopril, quinapril, ramipril, spirapril, temocapril, and trandolapril, and mixtures thereof.
25. The method according to Claim 19 wherein said anti-pressor agent is an  $\alpha_1$ -adrenergic antagonist selected from the group consisting of alfuzosin, apraclonidine, bunazosin, carvedilol, clonidine, dapiprazole, doxazosin, indoramin, labetalol, midrodrine, naphazoline, phenoxybenzamine, phentolamine, prazosin, tamsulosin, terazosin, trimazosin, and urapidil, and mixtures thereof.
26. The method according to Claim 19 wherein said anti-pressor agent is a calcium channel blocker selected from the group consisting of bepridil, diltiazem, mibrefadil, nicardipine, nifedipine, nimopidine, and verapamil, and mixtures thereof.



27. The method according to Claim 19 wherein said anti-pressor agent is a phosphodiesterase inhibitor selected from the group consisting of amrinone and sildenafil.
- 3 28. The method according to Claim 19 wherein said anti-pressor compound is co-administered with a diuretic agent.
29. The method according to Claim 19 wherein said anti-pressor compound is co-administered with prostaglandin- $E_1$ .
- 10 30. The method of Claim 18 wherein said anti-pressor agent is administered on a chronic basis at a dose ranging between one-twentieth to one-half the dose normally given to a hypertensive patient.
- 15 31. The method of Claim 18 wherein said anti-pressor agent is administered for a period ranging between about three days to about twenty-one days at a dose ranging between one to three times the dose normally given to a hypertensive patient.
32. The method according to Claim 30 wherein said anti-pressor agent is  
20 administered to a normotensive patient.
33. The method according to Claim 31 wherein said anti-pressor agent is administered to a normotensive patient.
- 25

1/6




Figure 1

2/6




Figure 2b

Figure 2a

3/6




Figure 2d




Figure 2c

4/6




Figure 3a

5/6




Figure 3b

6/6




Figure 4





**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>7</sup> :</b> <b>A61K 31/00, A61P 15/10</b>		<b>A3</b>	<b>(11) International Publication Number:</b> <b>WO 00/12110</b>
			<b>(43) International Publication Date:</b> 9 March 2000 (09.03.00)
<b>(21) International Application Number:</b> PCT/CA99/00787 <b>(22) International Filing Date:</b> 25 August 1999 (25.08.99) <b>(30) Priority Data:</b> 60/098,178                      26 August 1998 (26.08.98)                      US <b>(71) Applicant (for all designated States except US):</b> QUEEN'S UNIVERSITY AT KINGSTON [CA/CA]; Kingston, Ontario K7L 3N6 (CA). <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> ADAMS, Michael, A. [CA/CA]; 341 Renda Street, Kingston, Ontario K7M 5X9 (CA). HEATON, Jeremy, P., W. [CA/CA]; 1818 MacAlpine Road, R.R. #3, Gananoque, Ontario K7G 2V5 (CA). <b>(74) Agents:</b> STEEG, Carol, Miernicki et al.; Parteq Innovations, Queen's University at Kingston, Kingston, Ontario K7L 3N6 (CA).		<b>(81) Designated States:</b> AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i>  <b>(88) Date of publication of the international search report:</b> 3 August 2000 (03.08.00)	
<b>(54) Title:</b> USE OF ANTI(W)PRESSOR AGENTS FOR VASCULAR REMODELING IN GENITAL DYSFUNCTION			
<b>(57) Abstract</b> <p>The present invention provides a method of administration of an agent which acts to remodel neuronal or vascular pathways for the long term management of sexual dysfunction in both males and females. In a preferred embodiment, the invention provides a method of ameliorating or reversing pathogenic vascular degradative modeling in the ilio-hypogastric-pudental arterial bed and genitalia comprising administering to a human patient in need of such treatment a therapeutically effective amount of an anti-pressor agent. The anti-pressor agent comprises one or more compounds selected from the therapeutic classes of direct vasodilators such as hydralazine and NO donors, ACE inhibitors, angiotensin-II receptor antagonists, <math>\alpha_1</math>-adrenergic receptor antagonists, <math>\beta</math>-adrenergic receptor antagonists, calcium channel blockers, and phosphodiesterase inhibitors. The anti-pressor agent may be co-administered with a diuretic compound, and is administered either chronically at low dose, or for short periods of time at doses higher than are typically used for the treatment of hypertension. In certain embodiments of the method of the invention, the anti-pressor agent is co-administered with a diuretic agent and/or prostaglandin-E<sub>1</sub>.</p>			

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

# INTERNATIONAL SEARCH REPORT

In International Application No

PCT/CA 99/00787

**A. CLASSIFICATION OF SUBJECT MATTER**  
IPC 7 A61K31/00 A61P15/10

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61P A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 96 28142 A (VIVUS INC) 19 September 1996 (1996-09-19) page 3, line 30 -page 4, line 10 page 6, line 19 -page 7, line 23; claims	1-33
X	WO 97 33608 A (UNIV KINGSTON ;BANTING JAMES D (CA); HEATON JEREMY P W (CA); ADAMS) 18 September 1997 (1997-09-18) page 8, line 28 -page 9, line 3; claims	1-33
P,X	WO 98 58633 A (HEATON JEREMY P W ;UNIV KINGSTON (CA); ADAMS MICHAEL A (CA); BANTI) 30 December 1998 (1998-12-30) page 6, line 18 -page 7, line 27 page 9, line 1-7; claims	1-33
	-/-	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

\* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

15 May 2000

Date of mailing of the international search report

29/05/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Orviz Diaz, P

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/CA 99/00787

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	M. SCHARTL: "Remodeling of myocardium and arteries by chronic angiotensin converting enzyme inhibition in hypertensive patients" FILE BIOSIS, AN=PREV199497466956, XP002137640 abstract & J. HYPERTENSION, 12(SUPPL. 4):S37-S42, 1994,	1-33
A	R.M.K.W. LEE: "Debate: vascular remodeling" J. HYPERTENSION, vol. 15, no. 4, 1997, pages 333-337, XP000905246 cited in the application	1-33

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/CA 99/00787

### Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: —  
because they relate to subject matter not required to be searched by this Authority, namely:  
**Remark: Although all claims are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.**
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

### Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/CA 99/00787

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9628142 A	19-09-1996	AU 5358896 A	02-10-1996
		BR 9607365 A	30-12-1997
		CA 2215307 A	19-09-1996
		CZ 9702784 A	15-04-1998
		DE 814775 T	25-06-1998
		EP 0814775 A	07-01-1998
		ES 2110380 T	16-02-1998
		HU 9801324 A	28-09-1998
		JP 10505611 T	02-06-1998
		NO 974062 A	15-09-1997
		NZ 305560 A	25-11-1998
		PL 322268 A	19-01-1998
		SK 124397 A	02-12-1998
		US 5820587 A	13-10-1998
WO 9733608 A	18-09-1997	US 5688499 A	18-11-1997
		AU 2088597 A	01-10-1997
		CA 2248047 A	18-09-1997
		EP 0888123 A	07-01-1999
WO 9858633 A	30-12-1998	AU 8096198 A	04-01-1999
		EP 0998274 A	10-05-2000